



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Synthesis of Bis-Heterocyclic 1H-Imidazole 3-Oxides from 3-Oxido-1H-imidazole-4-carbohydrazides**

Pieczonka, A M ; Mlostoń, G ; Heimgartner, H

**Abstract:** The reaction of 1H-imidazole-4-carbohydrazides 1, which are conveniently accessible by treatment of the corresponding esters with  $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ , with isothiocyanates in refluxing EtOH led to thiosemicarbazides (= hydrazinecarbothioamides) 4 in high yields (Scheme 2). Whereas 4 in boiling aqueous NaOH yielded 2,4-dihydro-3H-1,2,4-triazole-3-thiones 5, the reaction in concentrated  $\text{H}_2\text{SO}_4$  at room temperature gave 1,3,4-thiadiazol-2-amines 6. Similarly, the reaction of 1 with butyl isocyanate led to semicarbazides 7, which, under basic conditions, undergo cyclization to give 2,4-dihydro-3H-1,2,4-triazol-3-ones 8 (Scheme 3). Treatment of 1 with  $\text{Ac}_2\text{O}$  yielded the diacylhydrazine derivatives 9 exclusively, and the alternative isomerization of 1 to imidazol-2-ones was not observed (Scheme 4). It is important to note that, in all these transformations, the imidazole N-oxide residue is retained. Furthermore, it was shown that imidazole N-oxides bearing a 1,2,4-triazole-3-thione or 1,3,4-thiadiazol-2-amine moiety undergo the S-transfer reaction to give bis-heterocyclic 1H-imidazole-2-thiones 11 by treatment with 2,2,4,4-tetramethylcyclobutane-1,3-dithione (Scheme 5).

DOI: <https://doi.org/10.1002/hlca.201100470>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-61480>

Journal Article

Accepted Version

Originally published at:

Pieczonka, A M; Mlostoń, G; Heimgartner, H (2012). Synthesis of Bis-Heterocyclic 1H-Imidazole 3-Oxides from 3-Oxido-1H-imidazole-4-carbohydrazides. *Helvetica Chimica Acta*, 95(3):404-414.

DOI: <https://doi.org/10.1002/hlca.201100470>

M-265 / 21.11.2011

Prof. Dr. H. Heimgartner

Tel. 044 635 4282

Fax 044 635 6836

e-mail: heimgart@oci.uzh.ch

**3-Oxidoimidazole-4-carbohydrazides for the Synthesis of some  
Bis-heterocyclic Imidazole *N*-Oxides**

by **Adam Marek Pieczonka<sup>1)</sup>**, **Grzegorz Mlostoń\***

University of Łódź, Department of Organic and Applied Chemistry, Tamka 12, PL-91-

403 Łódź (phone: +48 42 6355761; fax: +48 42 6655162; e-mail:

gmloston@uni.lodz.pl)

and **Heinz Heimgartner\***

Organisch-chemisches Institut der Universität Zürich, Winterthurerstrasse 190,

CH-8057 Zürich (phone: +41 44 6354282; fax: +41 44 6356812; e-mail:

heimgart@oci.uzh.ch)

---

<sup>1)</sup> Part of the planned Ph.D. thesis of *A. M. P.*, University of Łódź.

The reaction of 1*H*-imidazole-4-carbohydrazides **1**, which are conveniently accessible by treatment of the corresponding esters with  $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ , with isothiocyanates in refluxing EtOH led to thiosemicarbazides **4** in high yields (*Scheme 2*). Whereas **4** in boiling aqueous NaOH yielded 1,2,4-triazole-5-thiones **5**, the reaction in conc.  $\text{H}_2\text{SO}_4$  at room temperature gave 5-amino-1,3,4-thiadiazoles **6**. Similarly, the reaction of **1** with butylisocyanate led to semicarbazides **7**, which under basic conditions cyclized to give 1,2,4-triazol-5-ones **3** (*Scheme 3*). Treatment of **1** with  $\text{Ac}_2\text{O}$  yielded the diacylhydrazines **9** exclusively, and the alternative isomerization of **1** to imidazol-2-ones was not observed (*Scheme 4*). It is important to note that in all these transformations the imidazole *N*-oxide residue is retained. Furthermore, it was shown that imidazole *N*-oxides bearing a 1,2,4-triazole-5-thione or 5-amino-1,3,4-thiadiazole substituent undergo the S-transfer reaction to give bis-heterocyclic imidazole-2-thiones **11** by treatment with 2,2,4,4-tetramethylcyclobutane-1,3-dithione (*Scheme 5*).

**1. Introduction.** - Acid hydrazides (carbohydrazides) and the corresponding hydrazones are well known as a class of compounds with diverse biological activities [1]. Particularly important are derivatives bearing a heterocycle, and one of the best-known examples is isoniazid (isonicotinohydrazide) [2]. Imidazole derived hydrazides were also reported to display biological activity [3], and the majority of reports remain to patents (*e.g.*, ref. [4]). However, in most cases, *N,N*-disubstituted hydrazides were described, which are of limited interest with respect to their use as building blocks for new heterocyclic systems.

The *N*-unsubstituted carbohydrazides are widely applied as unique building blocks for the synthesis of diverse five- and six-membered heterocycles such as pyrroles, pyrazoles, thiazolidines, 1,2,4-triazoles, phthalazines, pyrimidines, etc. [5]. A typical procedure for the preparation of unsubstituted hydrazides is the treatment of the corresponding carboxylic ester with  $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$  [1][5]. In a recent publication we have reported on the application of this method for the preparation of diverse types of unsubstituted hydrazides containing in their structure the imidazole ring [6]. The precursors of 4-imidazolyl hydrazides **1** were the esters **2**, available *via* condensation of ethyl 2-hydroxyimino-3-oxobutanoate (**3**), a primary amine, and formaldehyde [7] (*Scheme 1*).

#### *Scheme 1*

In earlier papers, we presented the synthetic potential of the ‘nitron-like’ fragment of 2-unsubstituted imidazole *N*-oxides [8]. The goal of the present study was the synthesis of new bis-heterocycles with the preserved imidazole *N*-oxide unit by exploration of the reactivity of the carbohydrazido group in compounds **1**. In

preliminary experiments it was shown that hydrazides **1** can be used for the preparation of bis-heterocycles [6]. To the best of our knowledge, there are no other bis-heterocycles with the reactive *N*-oxide function known to date (cf. [6]).

**Results and Discussion.** – As described earlier, imidazole *N*-oxides **1** with the hydrazido substituent of C(4) are stable, colorless, crystalline materials [6]. In contrast to the corresponding esters they did not undergo the thermal isomerization to imidazole-2-ones. Even more surprisingly, they could neither be converted to the corresponding imidazole-2-thiones *via* the ‘sulfur-transfer reaction’ [8a] nor reduced to the parent system by treatment with *Raney*-Ni [8c].

It seems likely that these diminished reactivity of the N→O function is the result of a strong intramolecular H-bond [6][8d]. On the other hand, reactions with aldehydes and aliphatic ketones occurred smoothly yielding the expected hydrazones [6]. In addition, a preliminary experiment showed that thiosemicarbazide **4a** obtained from **1a** (R = PhCH<sub>2</sub>) and methyl isothiocyanate underwent the cyclocondensation yielding, depending on the reaction conditions, bis-heterocycles **5a** or **6a**. In analogy to **1a**, derivatives **1b** and **1c** were also reacted with other isothiocyanates (R<sup>2</sup> = Bu, Ph), and the thiosemicarbazides **4** were heated in a 2% aqueous solution of NaOH to give the triazole-3-thiones **5** (*Scheme 2*). In addition, the sterically crowded *tert*-butylisothiocyanate was used for the reaction with **1a**, and the expected thiosemicarbazide **4d** was obtained in good yield as a crystalline material. However, the attempted cyclization of **4d** under basic conditions was unsuccessful and the starting material was recovered in nearly quantitative yield. Apparently, in this case, steric hindrance resulting from the presence of the bulky *tert*-butyl group at the N-atom does not allow the system to undergo the cyclization.

## Scheme 2

The structure of the triazole-3-thiones **5** was indicated by the C=S absorption in the  $^{13}\text{C}$ -NMR spectra at 168 – 169 ppm. In addition, the unchanged *N*-oxide structure of the imidazole was confirmed by the characteristic  $^1\text{H}$ -NMR absorption of H–C(2) at 8.6 – 8.1 ppm.

The alternative course of the cyclocondensation was observed when thiosemicarbazides of type **4** were stirred in conc.  $\text{H}_2\text{SO}_4$  at room temperature for one day. Under these conditions, the only products formed were 5-amino-1,3,4-thiadiazol-2-yl-substituted imidazole *N*-oxides **6** (Scheme 2). Unexpectedly, the reactions carried out under these conditions with **4b** and **4i**, which were derived from phenylisothiocyanate, yielded **6b** and **6e**, respectively, bearing a phenylsulfonic acid residue ( $\text{R}^2 = 4\text{-HO}_3\text{SC}_6\text{H}_4$ ). It is very likely that these products result from the sulfonation of the intermediate 5-phenylamino-1,3,4-thiadiazole derivatives formed after the initial cyclocondensation step<sup>2</sup>). Based on the comparison of the chemical shifts of H–C(2) in **6b** and in the analogous MeN and BuN derivatives **6a** and **6c**, respectively, a zwitterionic structure can be postulated for both arylsulfonic acids **6b** and **6e** (see Scheme 5). To the best of our knowledge, secondary sulfonation of 5-phenylamino-1,3,4-thiadiazoles obtained from corresponding thiosemicarbazides in the presence of  $\text{H}_2\text{SO}_4$  under similar reaction conditions (r.t.) has not been reported so far [12][13].

---

<sup>2</sup>) The sulfonation of aniline with conc.  $\text{H}_2\text{SO}_4$  at elevated temperatures is a well known process [9] and has been studied in detail by *Khelevin* [10]. A study of the thermolysis of diphenylammonium hydrogen sulfate leading to 4-(phenylamino)benzenesulfonic acid has been published recently [11].

Similarly to the reactions with isothiocyanates, hydrazides **1a** and **1b** reacted with butyl isocyanate in refluxing EtOH yielding semicarbazides **7a** and **7b**, respectively, as crystalline materials in high yields (*Scheme 3*). After isolation and purification, they were used for the synthesis of bis-heterocyclic 1,2,4-triazol-3-ones **8a,b** under basic conditions. Both products **7** and **8** were obtained with the preserved *N*-oxide function.

### *Scheme 3*

Transformation of hydrazides into 3-alkyl-1,3,4-oxadiazoles can be performed *via* cyclization of their *N*-acyl derivatives under acidic conditions [5]. In the case of hydrazides **1a** and **1c**, the acetylation performed with a slight excess of Ac<sub>2</sub>O in boiling EtOH afforded the expected *N*-acetyl derivatives **9** (*Scheme 4*). Apparently, this conversion is much faster than the alternative isomerization to the corresponding imidazol-2-ones reported as a typical reaction of 2-unsubstituted imidazole *N*-oxides by treatment with Ac<sub>2</sub>O [8b]. However, the attempted cyclocondensations of products **9** under acidic conditions (H<sub>2</sub>SO<sub>4</sub>, r.t.) were unsuccessful.

### *Scheme 4*

In a series of reports we demonstrated that the treatment of 2-unsubstituted imidazole *N*-oxides with a cycloaliphatic thioketone, *e.g.*, 2,2,4,4-tetramethylcyclobutane-1,3-dithione (**10**) opens an easy access to imidazole-2-thiones [8a,c,e,f]. On the other hand, we found recently that the presence of a hydrazide group at C(4) inhibits the S-transfer reaction [6]. Having in hand bis-heterocyclic imidazole *N*-

oxides of type **5** and **6**, the S-transfer with **10** was tested. In both cases (**5g**, **6e**), the desired imidazole-2-thiones were smoothly formed at room temperature (*Scheme 5*).

#### *Scheme 5*

This result points out the influence of the intramolecular H-bond on the ‘nitronelike’ reactivity of the imidazole *N*-oxides **1**. Whereas the strong H-bond in the hydrazides [6] causes reduced reactivity, there is no such H-bond present in the bis-heterocyclic imidazole *N*-oxides **5** and **6**, which easily enter a [2 + 3] cycloaddition with **10**.

**Conclusions.** – The present study shows that imidazole carbohydrazides of type **1**, containing the *N*-oxide function, react with isothiocyanates, isocyanates, and Ac<sub>2</sub>O in a typical manner yielding the corresponding thiosemicarbazides **4**, semicarbazides **7**, and *N*-acetyl derivatives **9**, respectively, in high yields. In the case of thiosemicarbazides, cyclizations to 1,2,4-triazole-3-thiones **5** and 2-amino-1,3,4-thiadiazoles **6** were performed under basic and acidic conditions, correspondingly. Similarly, semicarbazides **7** were converted into 1,2,4-triazole-3-ones **8** in aqueous NaOH solution. In all these reactions, the *N*-oxide function in the imidazole ring was preserved. However, attempted acid-catalyzed cyclizations of **7** and **9** to the corresponding 3-amino- and 3-methyl-1,3,4-oxadiazoles, respectively, were unsuccessful.

In contrast to hydrazides **1**, which were shown not to undergo the S-transfer reaction upon treatment with the dithione **10** under standard conditions [6], bis-heterocyclic imidazole *N*-oxides of type **5** and **6** smoothly undergo this reaction.



To the best of our knowledge, the elaborated protocol offers a unique route for the synthesis of bis-heterocyclic *N*-oxides, which cannot be prepared using an oxidative approach. This novel type of *N*-oxides may be of interest as a new group of ligands for coordination chemistry. Moreover, in a recent paper, derivatives of 5-amino-1,3,4-thiadiazoles obtained from 5-methylimidazole-4-carbohydrazide according to the general method described in the present paper, displayed high anti-*Toxoplasma gondi* and antimicrobial activities [13].

The authors thank PD Dr. *L. Bigler*, University of Zurich, for recording of the HR-ESI-MS spectra. *A. P.* thanks for financial support within the project co-funded by the *European Union* under the *European Social Fund* ‘*HUMAN – BEST INVESTMENT!*’

## Experimental Part

*I. General.* M.p.: *STUART SMP30*; uncorrected. IR Spectra: *NEXUS FT-IR* spectrophotometer; in KBr; absorptions in  $\text{cm}^{-1}$ .  $^1\text{H}$ - and  $^{13}\text{C}\{^1\text{H}\}$ -NMR Spectra: *Bruker Avance III 600*, in  $(\text{D}_6)\text{DMSO}$ , using solvent signals as reference;  $\delta$  in ppm; coupling constants *J* in Hz; assignments of signals in  $^{13}\text{C}$ -NMR spectra accomplished on the basis of HMQC experiments. HR-ESI-MS: *Finnigan MAT-95*; HR-EI-MS: *Bruker Esquire LC* spectrometer.

2. *Starting Materials.* All solvents are commercially available and used as received. Hydrazides **1a** – **1c** were prepared from the corresponding esters by treatment with  $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$  according to the previously described protocol [6].

3. *Synthesis of Thiosemicarbazides 4a – 4j and Semicarbazides 7a and 7b.*

*General Procedure.* A mixture of hydrazide **1** (1 mmol) and the corresponding isothiocyanate (1.1 mmol) or isocyanate (1.1 mmol) in EtOH (5 ml) was heated to reflux for 2 h. Then, the formed product was filtered off, washed with  $\text{Et}_2\text{O}$  and crystallized from MeOH.

*1-[[[(1-Benzyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino]-3-methylthiourea (4a)* [6].

*1-[[[(1-Benzyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino]-3-phenylthiourea (4b):* Yield: 0.347 g (91%). Colorless crystals. M.p. 200 – 204° (dec., MeOH). IR (KBr): 3295s (NH), 3127s, 1663vs (C=O), 1599vs, 1497m, 708m.  $^1\text{H}$ -NMR ( $(\text{D}_6)$ DMSO): 9.80 (br. s, NH); 8.72 (s, H-C(2')); 7.48 – 7.12 (m, 10 arom. H); 5.26 (s,  $\text{CH}_2$ ); 2.45 (s, Me); two NH absorptions missing.  $^{13}\text{C}$ -NMR ( $(\text{D}_6)$ DMSO): 178.2 (C=O); 159.2 (C=S); 139.7, 135.9, 131.2, 121.3 (2 arom. C, C(4'), C(5')); 126.7 (C(2')); 129.5, 128.7, 128.6, 127.7, 127.6, 125.3 (10 arom. CH); 48.7 ( $\text{CH}_2$ ); 9.8 (Me). HR-ESI-MS: 404.1149 ( $[\text{M}+\text{Na}]^+$ ,  $\text{C}_{19}\text{H}_{19}\text{N}_5\text{NaO}_2\text{S}$ ; calc. 404.1152); 382.1329 ( $[\text{M}+\text{H}]^+$ ,  $\text{C}_{19}\text{H}_{20}\text{N}_5\text{O}_2\text{S}$ ; calc. 382.1332).

*1-[[[(1-Benzyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino]-3-butylthiourea (4c).* Yield: 0.343 g (95%). Colorless crystals. M.p. 204 – 207° (MeOH). IR (KBr): 3243m (NH), 2955s, 1661vs (C=O), 1598vs, 1552m, 1456m, 737m.  $^1\text{H}$ -NMR ( $(\text{D}_6)$ DMSO): 12.41, 9.31 (2 br. s, 2 NH); 8.72 (s, H-C(2')); 8.03 (br. s, NH); 7.42 – 7.25 (m, 5 arom. H); 5.25 (s,  $\text{CH}_2$ ); 3.40 (q,  $J = 6.6$ ,  $\text{CH}_2\text{N}$ ); 2.43 (s, Me); 1.49 – 1.44,

1.28 – 1.22 (2 *m*, 2 CH<sub>2</sub>); 0.86 (*t*, *J* = 7.2, Me). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 181.7 (C=O); 159.6 (C=S); 135.9, 131.2, 121.3 (1 arom. C, C(4'), C(5')); 126.7 (C(2')); 129.5, 128.7, 127.7 (5 arom. CH); 48.7 (PhCH<sub>2</sub>); 43.9 (CH<sub>2</sub>N); 31.3, 19.9 (2 CH<sub>2</sub>); 14.2 (Me); 9.8 (MeC(2')). HR-ESI-MS: 384.1467 ([*M*+Na]<sup>+</sup>, C<sub>17</sub>H<sub>23</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 384.1465); 362.1643 ([*M*+H]<sup>+</sup>, C<sub>17</sub>H<sub>24</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 362.1645).

*1-[(1-Benzyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino-3-(tert-butyl)thiourea (4d)*. Yield: 0.235 g (65%). Colorless crystals. M.p. 201 – 203° (MeOH). IR (KBr): 3235*m* (NH), 3088*s*, 1668*vs* (C=O), 1599*vs*, 1541*m*, 1362*m*, 709*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 9.51 (br. *s*, NH); 8.67 (*s*, H–C(2')); 7.41 – 7.25 (*m*, 5 arom. H); 5.24 (*s*, CH<sub>2</sub>); 2.43 (*s*, Me); 1.44 (*s*, Me<sub>3</sub>C); two NH absorptions missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 178.9 (C=O); 159.8 (C=S); 135.8, 131.1, 120.9 (1 arom. C, C(4'), C(5')); 126.7 (C(2')); 129.4, 128.6, 127.7 (5 arom. CH); 53.2 (Me<sub>3</sub>C); 48.7 (CH<sub>2</sub>); 29.1 (3 Me<sub>3</sub>C); 9.8 (MeC(2')). HR-ESI-MS: 384.1469 ([*M*+Na]<sup>+</sup>, C<sub>17</sub>H<sub>23</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 384.1465); 362.1646 ([*M*+H]<sup>+</sup>, C<sub>17</sub>H<sub>24</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 362.1645).

*1-[(1,5-Dimethyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino-3-methylthiourea (4e)*. Yield: 0.173 g (71%). Colorless crystals. M.p. 229 – 231° (MeOH). IR (KBr): 3160*s*, 2974*s*, 1638*vs* (C=O), 1602*vs*, 1485*s*, 1285*m*, 606*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 9.35 (br. *s*, NH); 8.48 (*s*, H–C(2')); 8.01 (br. *s*, NH); 3.58 (*s*, imidazole Me); 2.85 (*d*, *J* = 4.1, MeN); 2.48 (*s*, Me); one NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 182.6 (C=O); 159.8 (C=S); 131.8, 120.7 (C(4'), C(5')); 126.8 (C(2')); 32.4 (MeN(1')); 31.4 (MeN); 9.5 (MeC(2')). HR-ESI-MS: 266.0682 ([*M*+Na]<sup>+</sup>, C<sub>8</sub>H<sub>13</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 266.0682); 244.0859 ([*M*+H]<sup>+</sup>, C<sub>8</sub>H<sub>14</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 244.0863).

*1-[(1,5-Dimethyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino-3-phenylthiourea (4f)*. Yield: 0.262 g (86%). Colorless crystals. M.p. 223 – 225° (MeOH). IR (KBr): 3245*m* (NH), 3093*s*, 3036*s*, 1667*vs* (C=O), 1600*vs*, 1497*s*, 1321*s*, 760*m*. <sup>1</sup>H-

NMR ((D<sub>6</sub>)DMSO): 9.79 (br. *s*, NH); 8.48 (*s*, H–C(2')); 7.50 – 7.11 (*m*, 5 arom. H); 3.58 (*s*, MeN); 2.50 (*s*, Me; overlaps with DMSO signal); two NH absorptions missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 179.0 (C=O); 160.8 (C=S); 139.8, 131.6, 120.7 (1 arom. C, C(4'), C(5')); 128.8, 128.6, 125.1 (5 arom. CH); 126.8 (C(2')); 32.4 (MeN); 9.5 (MeC(2')). HR-ESI-MS: 328.0835 ([*M*+Na]<sup>+</sup>, C<sub>13</sub>H<sub>15</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 328.0839); 306.1013 ([*M*+H]<sup>+</sup>, C<sub>13</sub>H<sub>16</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 306.1019).

*1-[(1,5-Dimethyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino*-3-*butylthiourea* (**4g**). Yield: 0.262 g (92%). Colorless crystals. M.p. 208 – 210° (MeOH). IR (KBr): 3189*s*, 2957*s*, 1640*vs* (C=O), 1599*vs*, 1483*s*, 1277*m*, 616*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 12.44, 9.30 (2 br. *s*, 2 NH); 8.47 (*s*, H–C(2')); 8.01 (br. *s*, NH); 3.58 (*s*, MeN); 3.40 (*q*, *J* = 6.6, CH<sub>2</sub>N); 2.48 (*s*, Me); 1.49 – 1.44, 1.29 – 1.22 (2 *m*, 2 CH<sub>2</sub>); 0.87 (*t*, *J* = 7.2, Me). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 181.7 (C=O); 159.5 (C=S); 131.7, 120.7 (C(4'), C(5')); 126.8 (C(2')); 43.9 (CH<sub>2</sub>N); 31.4 (MeN); 31.3, 19.9 (2 CH<sub>2</sub>); 14.2 (Me); 9.5 (MeC(2')). HR-ESI-MS: 308.1151 ([*M*+Na]<sup>+</sup>, C<sub>11</sub>H<sub>19</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 308.1152); 286.1329 ([*M*+H]<sup>+</sup>, C<sub>11</sub>H<sub>20</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 286.1332).

*1-[(1-Cyclohexyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino*-3-*methylthiourea* (**4h**). Yield: 0.258 g (83%). Colorless crystals. M.p. 207 – 211° (MeOH). IR (KBr): 3250*s* (NH), 3139*s*, 1664*vs* (C=O), 1596*vs*, 1552*m*, 1495*m*, 1259*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 12.45, 9.35 (2 br. *s*, 2 NH); 8.72 (*s*, H–C(2')); 8.00 (br. *s*, NH); 4.12 – 4.07 (*m*, CH); 2.85 (*d*, *J* = 4.4, MeN); 2.55 (*s*, Me); 1.92 – 1.79 (*m*, 4 cyclohexyl H); 1.67 – 1.60 (*m*, 3 cyclohexyl H); 1.44 – 1.38 (*m*, 2 cyclohexyl H); 1.21 – 1.13 (*m*, 1 cyclohexyl H). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 182.6 (C=O); 159.9 (C=S); 130.7, 120.3 (C(4'), C(5')); 124.5 (C(2')); 55.1 (CH); 33.0, 25.3, 25.0 (5 cyclohexyl CH<sub>2</sub>); 31.4 (MeN); 9.5 (MeC(2')). HR-ESI-MS: 334.1306 ([*M*+Na]<sup>+</sup>, C<sub>13</sub>H<sub>21</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 334.1308); 312.1488 ([*M*+H]<sup>+</sup>, C<sub>13</sub>H<sub>22</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 312.1489).

*1-[(1-Cyclohexyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino*}-3-phenylthiourea (**4i**). Yield: 0.332 g (89%). Colorless crystals. M.p. 174 – 176° (MeOH). IR (KBr): 3231<sub>s</sub> (NH), 3139<sub>s</sub>, 2934<sub>s</sub>, 1675<sub>vs</sub> (C=O), 1600<sub>vs</sub>, 1497<sub>m</sub>, 1257<sub>m</sub>. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 9.81 (br. *s*, NH); 8.73 (*s*, H-C(2')); 7.50 – 7.12 (*m*, 5 arom. H); 4.12 – 4.07 (*m*, CH); 2.57 (*s*, Me); 1.93 – 1.80 (*m*, 4 cyclohexyl H); 1.67 – 1.61 (*m*, 3 cyclohexyl H); 1.45 – 1.38 (*m*, 2 cyclohexyl H); 1.22 – 1.14 (*m*, 1 cyclohexyl H); two NH absorptions missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 172.4 (C=O); 156.2 (C=S); 139.7, 130.6, 120.3 (1 arom. C, C(4'), C(5')); 129.4, 129.1, 128.6 (5 arom. CH); 124.4 (C(2')); 55.1 (CH); 33.0, 25.4, 25.0 (5 cyclohexyl CH<sub>2</sub>); 9.5 (MeC(2')). HR-ESI-MS: 396.1469 ([*M*+Na]<sup>+</sup>, C<sub>18</sub>H<sub>23</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 396.1465); 374.1649 ([*M*+H]<sup>+</sup>, C<sub>18</sub>H<sub>24</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 374.1645).

*1-[(1-Cyclohexyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino*}-3-butylthiourea (**4j**). Yield: 0.318 g (90%). Colorless crystals. M.p. 176 – 180° (MeOH). IR (KBr): 3302<sub>s</sub> (NH), 3143<sub>s</sub>, 2934<sub>s</sub>, 1681<sub>vs</sub> (C=O), 1604<sub>vs</sub>, 1545<sub>m</sub>, 1265<sub>m</sub>. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 12.48, 9.26 (2 br. *s*, 2 NH); 8.72 (*s*, H-C(2')); 8.02 (br. *s*, NH); 4.11 – 4.07 (*m*, CH); 3.43 – 3.38 (*m*, CH<sub>2</sub>N); 2.55 (*s*, Me); 1.92 – 1.79 (*m*, 4 cyclohexyl H); 1.67 – 1.60 (*m*, 3 cyclohexyl H); 1.49 – 1.38 (*m*, 2 cyclohexyl H, butyl CH<sub>2</sub>); 1.28 – 1.15 (*m*, 1 cyclohexyl H, butyl CH<sub>2</sub>); 0.87 (*t*, *J* = 7.6, Me). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 181.7 (C=O); 159.8 (C=S); 130.6, 120.3 (C(4'), C(5')); 124.4 (C(2')); 55.0 (CH); 43.9 (CH<sub>2</sub>N); 33.0, 25.3, 25.0 (5 cyclohexyl CH<sub>2</sub>); 31.3, 19.9 (2 butyl CH<sub>2</sub>); 14.2 (Me); 9.5 (MeC(2')). HR-ESI-MS: 376.1777 ([*M*+Na]<sup>+</sup>, C<sub>16</sub>H<sub>27</sub>N<sub>5</sub>NaO<sub>2</sub>S; calc. 376.1778); 354.1955 ([*M*+H]<sup>+</sup>, C<sub>16</sub>H<sub>28</sub>N<sub>5</sub>O<sub>2</sub>S; calc. 354.1958).

*1-[(1-Benzyl-5-methyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino*}-3-butylthiourea (**7a**). Yield: 0.338 g (98%). Colorless crystals. M.p. 201 – 205° (MeOH). IR (KBr): 3304<sub>s</sub> (NH), 3105<sub>m</sub>, 2960<sub>m</sub>, 1656<sub>vs</sub> (C=O), 1603<sub>s</sub>, 1544<sub>m</sub>. <sup>1</sup>H-NMR

((D<sub>6</sub>)DMSO): 12.08 (br. s, NH); 8.67 (s, H-C(2')); 7.41 – 7.23 (m, 5 arom. H); 7.94, 6.42 (2 br. s, 2 NH); 5.24 (s, CH<sub>2</sub>); 3.01 (q, *J* = 7.0, CH<sub>2</sub>N); 2.42 (s, Me); 1.38 – 1.34, 1.29 – 1.22 (2 m, 2 CH<sub>2</sub>); 0.86 (t, *J* = 7.0, Me). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 159.5, 158.0 (2 C=O); 135.9, 131.0, 121.3 (1 arom. C, C(4'), C(5')); 126.8 (C(2')); 129.4, 128.6, 127.6 (5 arom. CH); 48.7 (PhCH<sub>2</sub>); 39.4 (CH<sub>2</sub>N); 32.4, 19.9 (2 CH<sub>2</sub>); 14.2 (Me); 9.7 (MeC(2')). HR-ESI-MS: 368.1697 ([*M*+Na]<sup>+</sup>, C<sub>17</sub>H<sub>23</sub>N<sub>5</sub>NaO<sub>3</sub>; calc. 368.1693); 346.1875 ([*M*+H]<sup>+</sup>, C<sub>17</sub>H<sub>24</sub>N<sub>5</sub>O<sub>3</sub>; calc. 346.1874).

*1-[(1,5-Dimethyl-3-oxido-1H-imidazolium-4-yl)carbonyl]amino-3-butylurea* (**7b**). Yield: 0.256 g (95%). Colorless crystals. M.p. 190 – 193° (MeOH). IR (KBr): 3308vs, 3132m, 2957s, 1644vs (C=O), 1602s, 1534s, 1254m, 608m. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 12.10 (br. s, NH); 8.44 (s, H-C(2')); 7.93, 6.39 (2 br. s, 2 NH); 3.56 (s, MeN); 3.00 (q, *J* = 5.8, CH<sub>2</sub>N); 2.47 (s, Me); 1.39 – 1.34, 1.29 – 1.23 (2 m, 2 CH<sub>2</sub>); 0.87 (t, *J* = 7.0, Me). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 159.5, 158.0 (2 C=O); 131.4, 120.7 (C(4'), C(5')); 126.8 (C(2')); 39.4 (CH<sub>2</sub>N); 32.4 (MeN); 32.3, 19.9 (2 CH<sub>2</sub>); 14.2 (Me); 9.4 (MeC(2')). HR-ESI-MS: 292.1383 ([*M*+Na]<sup>+</sup>, C<sub>11</sub>H<sub>19</sub>N<sub>5</sub>NaO<sub>3</sub>; calc. 292.1380); 270.1560 ([*M*+H]<sup>+</sup>, C<sub>11</sub>H<sub>20</sub>N<sub>5</sub>O<sub>3</sub>; calc. 270.1561).

4. *Synthesis of 1,2,4-Triazole-3-thiones 5. General Procedure.* A mixture of thiosemicarbazide **4** (1 mmol) and a 2% aq. soln. of NaOH (5 ml) was heated to reflux for 2 h. Then, the soln. was neutralized with AcOH and the formed precipitate was filtered off and crystallized from MeOH.

*1-Benzyl-4-(4,5-dihydro-4-methyl-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide (5a)* [6].

*1-Benzyl-4-(4,5-dihydro-4-phenyl-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide (5b)*. Yield: 0.192 g (53%). Colorless crystals. M.p. 262 – 264° (dec. MeOH). IR (KBr): 3110s, 3044m, 1497s, 1319m, 688m. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.35 (s,

H–C(2)); 7.47 – 7.28 (*m*, 10 arom. H); 5.11 (*s*, CH<sub>2</sub>N); 1.92 (*s*, Me); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 168.8 (C=S); 141.4 (triazole C(3)); 134.3, 128.8, 136.2, 117.6 (2 arom. C, imidazole C(4), C(5)); 129.3, 129.2, 128.4, 127.7, 127.2, 126.8 (10 arom. CH); 126.2 (imidazole C(2)); 48.8 (CH<sub>2</sub>N); 8.9 (Me). HR-ESI-MS: 386.1041 ([*M*+Na]<sup>+</sup>, C<sub>19</sub>H<sub>17</sub>N<sub>5</sub>NaOS; calc. 386.1046); 364.1221 ([*M*+H]<sup>+</sup>, C<sub>19</sub>H<sub>18</sub>N<sub>5</sub>OS; calc. 364.1227).

*1-Benzyl-4-(4,5-dihydro-4-butyl-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide (5c)*. Yield: 0.240 g (70%). Colorless crystals. M.p. 256–258° (MeOH). IR (KBr): 3129*m*, 2930*m*, 1455*m*, 1294*m*, 730*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.60 (*s*, H–C(2)); 7.41 – 7.25 (*m*, 5 arom. H); 5.26 (*s*, CH<sub>2</sub>N); 4.12 (*t*, *J* = 6.8, CH<sub>2</sub>N); 2.08 (*s*, Me); 1.45 – 1.40 (*m*, CH<sub>2</sub>); 1.04 – 0.98 (*m*, CH<sub>2</sub>); 0.67 (*t*, *J* = 6.8, Me); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 167.9 (C=S); 141.7 (triazole C(3)); 136.1, 128.4, 118.5 (1 arom. C, imidazole C(4), C(5)); 129.4, 128.6, 127.6 (5 arom. CH); 126.7 (imidazole C(2)); 49.2 (CH<sub>2</sub>N); 44.0 (butyl CH<sub>2</sub>N); 30.1, 19.4 (2 butyl CH<sub>2</sub>); 13.7 (butyl Me); 9.1 (Me). HR-ESI-MS: 366.1359 ([*M*+Na]<sup>+</sup>, C<sub>17</sub>H<sub>21</sub>N<sub>5</sub>NaOS; calc. 366.1359); 344.1537 ([*M*+H]<sup>+</sup>, C<sub>17</sub>H<sub>22</sub>N<sub>5</sub>OS; calc. 344.1540).

*1,5-Dimethyl-4-(4,5-dihydro-4-methyl-5-thioxo-1H-1,2,4-triazol-3-yl)-1H-imidazole 3-Oxide (5d)*. Yield: 0.144 g (64%). Colorless crystals. M.p. 321 – 323° (MeOH). IR (KBr): 3147*vs* (NH), 1515*m*, 1455*m*, 1329*m*, 929*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.31 (*s*, H–C(2)); 3.58, 3.44 2(*s*, 2 MeN); 2.17 (*s*, Me); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 168.2 (C=S); 142.3 (triazole C(3)); 128.1, 118.7 (imidazole C(4), C(5)); 126.2 (imidazole C(2)); 32.7, 31.4 (2 MeN); 9.1 (Me). HR-ESI-MS: 248.0574 ([*M*+Na]<sup>+</sup>, C<sub>8</sub>H<sub>11</sub>N<sub>5</sub>NaOS; calc. 248.0577); 226.0755 ([*M*+H]<sup>+</sup>, C<sub>8</sub>H<sub>12</sub>N<sub>5</sub>OS; calc. 226.0757).

*1,5-Dimethyl-4-(4,5-dihydro-4-phenyl-5-thioxo-1H-1,2,4-triazol-3-yl)-1H-imidazole 3-Oxide (5e)*. Yield: 0.155 g (54%). Colorless crystals. M.p. 270 – 274° (dec. MeOH). IR (KBr): 3156<sub>vs</sub> (NH), 1506<sub>m</sub>, 1497<sub>s</sub>, 1326<sub>m</sub>, 694<sub>m</sub>. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.14 (s, H–C(2)); 7.51 – 7.38 (m, 5 arom. H); 3.45 (s, MeN); 2.09 (s, Me); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 169.0 (C=S); 141.7 (triazole C(3)); 136.9, 128.9, 117.1 (1 arom. C, imidazole C(4), C(5)); 129.4, 129.1, 127.8 (5 arom. CH); 126.2 (imidazole C(2)); 32.6 (MeN); 8.7 (Me). HR-ESI-MS: 310.0733 ([M+Na]<sup>+</sup>, C<sub>13</sub>H<sub>13</sub>N<sub>5</sub>NaOS; calc. 310.0733); 288.0911 ([M+H]<sup>+</sup>, C<sub>13</sub>H<sub>14</sub>N<sub>5</sub>OS; calc. 288.0914).

*1,5-Dimethyl-4-(4-butyl-4,5-dihydro-5-thioxo-1H-1,2,4-triazol-3-yl)-1H-imidazole 3-Oxide (5f)*. Yield: 0.143 g (50%). Colorless crystals. M.p. 240 – 242° (MeOH). IR (KBr): 3162<sub>s</sub> (NH), 2953<sub>s</sub>, 1570<sub>m</sub>, 1449<sub>m</sub>, 1348<sub>m</sub>, 1295<sub>m</sub>, 926<sub>m</sub>. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.38 (s, H–C(2)); 4.14 (t, *J* = 7.7, CH<sub>2</sub>N); 3.59 (s, MeN); 2.17 (s, Me); 1.51 – 1.46, 1.12 – 1.05 (2<sub>m</sub>, 2 CH<sub>2</sub>); 0.74 (t, *J* = 7.7, Me); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 167.9 (C=S); 142.2 (triazole C(3)); 128.8, 117.5 (imidazole C(4), C(5)); 126.7 (imidazole C(2)); 44.1 (CH<sub>2</sub>N); 32.8 (MeN); 30.0, 19.5 (2 CH<sub>2</sub>); 13.7, 8.9 (2 Me). HR-ESI-MS: 290.1042 ([M+Na]<sup>+</sup>, C<sub>11</sub>H<sub>17</sub>N<sub>5</sub>NaOS; calc. 290.1046); 268.1225 ([M+H]<sup>+</sup>, C<sub>11</sub>H<sub>18</sub>N<sub>5</sub>OS; calc. 268.1227).

*1-Cyclohexyl-4-(4,5-dihydro-4-methyl-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide (5g)*. Yield: 0.240 g (82%). Colorless crystals. M.p. 212 – 216° (MeOH). IR (KBr): 3115<sub>s</sub> (NH), 2941<sub>s</sub>, 1559<sub>m</sub>, 1418<sub>m</sub>, 1324<sub>m</sub>. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.49 (s, H–C(2)); 4.05 – 4.00 (m, CH); 3.42 (s, MeN); 2.20 (s, Me); 1.97 – 1.80 (m, 4 cyclohexyl H); 1.69 – 1.62 (m, 3 cyclohexyl H); 1.45 – 1.38 (m, 2 cyclohexyl H); 1.23 – 1.14 (m, 1 cyclohexyl H); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 168.3 (C=S); 142.1 (triazole C(3)); 126.7, 118.7 (imidazole C(4), C(5)); 123.6 (imidazole C(2)); 55.5 (CH); 33.2 (MeN); 31.4, 25.4, 23.1 (5 cyclohexyl CH<sub>2</sub>); 9.2 (Me). HR-ESI-



MS: 316.1200 ( $[M+Na]^+$ ,  $C_{13}H_{19}N_5NaOS$ ; calc. 316.1203); 294.1379 ( $[M+H]^+$ ,  $C_{13}H_{20}N_5OS$ ; calc. 294.1383).

*1-Cyclohexyl-4-(4,5-dihydro-4-phenyl-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide (5h)*. Yield: 0.185 g (52%). Colorless crystals. M.p. 282 – 284° (MeOH). IR (KBr): 3231m (NH), 2936m, 1601m, 1497m, 1320m.  $^1H$ -NMR ( $(D_6)$ DMSO): 8.34 (s, H-C(2)); 7.45 – 7.37 (m, 5 arom. H); 3.92 – 3.85 (m, CH); 2.07 (s, Me); 1.79 – 1.74 (m, 4 cyclohexyl H); 1.62 – 1.49 (m, 3 cyclohexyl H); 1.37 – 1.30 (m, 2 cyclohexyl H); 1.15 – 1.12 (m, 1 cyclohexyl H); NH absorption missing.  $^{13}C$ -NMR ( $(D_6)$ DMSO): 168.9 (C=S); 141.6 (triazole C(3)); 134.1, 126.1, 118.4 (1 arom. C, imidazole C(4), C(5)); 129.3, 129.0, 127.7 (5 arom. CH); 123.6 (imidazole C(2)); 55.4 (CH); 33.0, 25.3, 24.9 (5 cyclohexyl  $CH_2$ ); 8.9 (Me). HR-ESI-MS: 378.1361 ( $[M+Na]^+$ ,  $C_{18}H_{21}N_5NaOS$ ; calc. 378.1359); 356.1541 ( $[M+H]^+$ ,  $C_{18}H_{22}N_5OS$ ; calc. 356.1540).

*1-Cyclohexyl-4-(4-butyl-4,5-dihydro-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide (5i)*. Yield: 0.251 g (75%). Colorless crystals. M.p. 248–250° (MeOH). IR (KBr): 3066m, 2936m, 1356m, 1295m.  $^1H$ -NMR ( $(D_6)$ DMSO): 8.55 (s, H-C(2)); 4.12 (t,  $J = 6.8$ ,  $CH_2N$ ); 4.07 – 4.02 (m, CH); 2.20 (s, Me); 1.95 – 1.81 (m, 4 cyclohexyl H); 1.69 – 1.62 (m, 3 cyclohexyl H); 1.48 – 1.38 (m, 2 cyclohexyl H, butyl  $CH_2$ ); 1.22 – 1.17 (m, 1 cyclohexyl H); 1.07 – 1.01 (m, butyl  $CH_2$ ); 0.69 (t,  $J = 6.8$ , Me); NH absorption missing.  $^{13}C$ -NMR ( $(D_6)$ DMSO): 167.9 (C=S); 141.9 (triazole C(3)); 123.1, 118.4 (imidazole C(4), C(5)); 123.9 (imidazole C(2)); 55.6 (CH); 43.7 ( $CH_2N$ ); 33.2, 25.4, 25.0 (5 cyclohexyl  $CH_2$ ); 30.0, 19.3 (2 butyl  $CH_2$ ); 13.6, 9.0 (2 Me). HR-ESI-MS: 358.1677 ( $[M+Na]^+$ ,  $C_{16}H_{25}N_5NaOS$ ; calc. 358.1672); 336.1856 ( $[M+H]^+$ ,  $C_{16}H_{26}N_5OS$ ; calc. 336.1853).

5. *Synthesis of 1,3,4-Thiadiazoles* 6. *General procedure*. A soln. of thiosemicarbazide **4** (1 mmol) in conc.  $H_2SO_4$  (5 ml) was kept at r.t. for 1 d. After

neutralization of the soln. with diluted  $\text{NH}_4\text{OH}$ , the solid product was filtered off, dried, and crystallized from MeOH.

*1-Benzyl-5-methyl-4-(5-methylamino-1,3,4-thiadiazol-2-yl)-1H-imidazole 3-Oxide (6a)* [6].

*1-Benzyl-5-methyl-4-[5-(4-sulfophenyl)amino]-1,3,4-thiadiazol-2-yl)-1H-imidazole 3-Oxide (6b)*. Yield: 0.211 g (58%). Yellowish crystals. M.p.  $300 - 303^\circ$  (dec., MeOH). IR (KBr):  $3400-2800m$  (br),  $3124m$ ,  $3064m$ ,  $1509s$ ,  $1176m$  (br),  $1034m$ ,  $708m$ .  $^1\text{H-NMR}$  ( $(\text{D}_6)$ DMSO): 10.34 (br. s, NH); 8.66 (s, H-C(2)); 7.60, 7.57 (AA'BB',  $J_{\text{AB}} = 8.4$ , 4 arom. H); 7.42 – 7.28 (m, 5 arom. H); 5.29 (s,  $\text{CH}_2$ ); 2.56 (s, Me).  $^{13}\text{C-NMR}$  ( $(\text{D}_6)$ DMSO): 164.4, 146.4 (thiadiazole C(2), C(5)); 142.1, 141.4, 136.1, 122.7, 116.6 (3 arom. C, imidazole C(4), C(5)); 129.4, 128.6, 127.7, 127.0, 125.0 (9 arom. CH); 125.6 (imidazole C(2)); 49.0 ( $\text{CH}_2$ ); 10.2 (Me). HR-ESI-MS: 488.0432 ( $[\text{M}-1+2 \text{ Na}]^+$ ,  $\text{C}_{19}\text{H}_{16}\text{N}_5\text{Na}_2\text{O}_4\text{S}_2$ ; calc. 488.0434); 466.0610 ( $[\text{M}+\text{Na}]^+$ ,  $\text{C}_{19}\text{H}_{17}\text{N}_5\text{NaO}_4\text{S}_2$ ; calc. 466.0614); 444.0791 ( $[\text{M}+\text{H}]^+$ ,  $\text{C}_{19}\text{H}_{18}\text{N}_5\text{O}_4\text{S}_2$ ; calc. 444.0795).

*1-Benzyl-5-methyl-4-(5-butylamino-1,3,4-thiadiazol-2-yl)-1H-imidazole 3-Oxide (6c)*. Yield: 0.209 g (61%). Yellowish crystals. M.p.  $206 - 208^\circ$  (dec., MeOH). IR (KBr):  $3185m$ ,  $3068m$ ,  $2958s$ ,  $1576s$ ,  $1454m$ ,  $748m$ .  $^1\text{H-NMR}$  ( $(\text{D}_6)$ DMSO): 8.57 (s, H-C(2)); 7.58 (br. s, NH); 7.41 – 7.26 (m, 5 arom. H); 5.24 (s,  $\text{PhCH}_2$ ); 3.34 – 3.23 (m,  $\text{CH}_2\text{N}$ ); 2.49 (s, Me); 1.59 – 1.54 (m,  $\text{CH}_2$ ); 1.39 – 1.33 (m,  $\text{CH}_2$ ); 0.90 (t,  $J = 7.6$ , Me).  $^{13}\text{C-NMR}$  ( $(\text{D}_6)$ DMSO): 169.2, 144.4 (thiadiazole C(2), C(5)); 136.3, 124.0, 123.0 (1 arom. C, imidazole C(4), C(5)); 129.4, 128.5, 127.6 (5 arom. CH); 125.3 (imidazole C(2)); 48.9 ( $\text{PhCH}_2$ ); 44.9 ( $\text{CH}_2\text{N}$ ); 31.2, 20.1 (2  $\text{CH}_2$ ); 14.1, 10.1 (2 Me). HR-ESI-MS: 366.1361 ( $[\text{M}+\text{Na}]^+$ ,  $\text{C}_{17}\text{H}_{21}\text{N}_5\text{NaOS}$ ; calc. 366.1359); 344.1541 ( $[\text{M}+\text{H}]^+$ ,  $\text{C}_{17}\text{H}_{22}\text{N}_5\text{OS}$ ; calc. 344.1540).

*1,5-Dimethyl-4-(5-methylamino-1,3,4-thiadiazol-2-yl)-1H-imidazole 3-Oxide* (**6d**). Yield: 0.207 g (92%). Yellowish crystals. M.p. 228 – 230° (dec., MeOH). IR (KBr): 3375s (NH), 3120s, 2941s, 1541m, 1404m, 1097m, 1033m, 604m. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 8.37 (s, H–C(2)); 7.55 (br. s, NH); 3.59 (s, imidazole MeN); 2.89 (br. s, MeN); 2.54 (s, Me). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 169.8, 144.8 (thiadiazole C(2), C(5)); 125.4 (imidazole C(2)); 124.5, 122.3 (imidazole C(4), C(5)); 32.5, 31.6 (2 MeN); 9.8 (Me). HR-ESI-MS: 248.0578 ([M+Na]<sup>+</sup>, C<sub>8</sub>H<sub>11</sub>N<sub>5</sub>NaOS; calc. 248.0577); 226.0760 ([M+H]<sup>+</sup>, C<sub>8</sub>H<sub>12</sub>N<sub>5</sub>OS; calc. 226.0757).

*1-Cyclohexyl-5-methyl-4-[5-(4-sulfophenyl)amino]-1,3,4-thiadiazol-2-yl)-1H-imidazole 3-Oxide* (**6e**). Yield: 0.249 g (70%). Colorless crystals. M.p. 310 – 312° (dec., MeOH). IR (KBr): 3400–2650m (br), 3385s (NH), 3227s, 3038s, 1420s (br), 1090s (br), 613m. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 10.29 (br. s, NH); 8.65 (s, H–C(2)); 7.60, 7.56 (AA'BB', *J*<sub>AB</sub> = 8.4, 4 arom. H); 4.14 – 4.09 (m, CH); 2.68 (s, Me); 1.99 – 1.81 (m, 4 cyclohexyl H); 1.68 – 1.63 (m, 3 cyclohexyl H); 1.47 – 1.41 (m, 2 cyclohexyl H); 1.24 – 1.21 (m, 1 cyclohexyl H). <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 164.4, 146.7 (thiadiazole C(2), C(5)); 142.2, 116.6 (2 arom. C); 141.4, 127.0 (4 arom. CH); 124.2 (imidazole C(2)); 123.1, 121.6 (imidazole C(4), C(5)); 55.3 (CH); 33.1, 25.4, 25.0 (5 cyclohexyl CH<sub>2</sub>); 9.9 (Me). HR-ESI-MS: 458.0926 ([M+Na]<sup>+</sup>, C<sub>18</sub>H<sub>21</sub>N<sub>5</sub>NaO<sub>4</sub>S<sub>2</sub>; calc. 458.0927); 436.1110 ([M+H]<sup>+</sup>, C<sub>18</sub>H<sub>22</sub>N<sub>5</sub>O<sub>4</sub>S<sub>2</sub>; calc. 436.1108).

6. *Synthesis of 1,2,4-Triazole-3-ones 8. General procedure.* A mixture of semicarbazide **7** (1 mmol) and a 2% aq. soln. of NaOH (5 ml) was heated to reflux for 8 h. Then, the soln. was neutralized with AcOH, the formed precipitate was filtered off, and crystallized from MeOH.

*1-Benzyl-4-(4-butyl-4,5-dihydro-5-oxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole 3-Oxide* (**8a**). Yield: 0.216 g (66%). Colorless crystals. M.p. 224 – 228°

(MeOH). IR (KBr): 3115 $m$ , 2938 $m$ , 1705 $vs$  (C=O), 1558 $m$ , 1416 $m$ , 734 $m$ .  $^1\text{H-NMR}$  ((D<sub>6</sub>)DMSO): 8.55 ( $s$ , H–C(2)); 7.41 – 7.24 ( $m$ , 5 arom. H); 5.23 ( $s$ , PhCH<sub>2</sub>); 3.73 ( $t$ ,  $J$  = 6.8, CH<sub>2</sub>N); 2.08 ( $s$ , Me); 1.58 (br.  $s$ , NH); 1.37 – 1.32, 1.06 – 1.00 ( $2m$ , 2 CH<sub>2</sub>); 0.68 ( $t$ ,  $J$  = 7.7, Me).  $^{13}\text{C-NMR}$  ((D<sub>6</sub>)DMSO): 155.4 (C=O); 137.4, 136.3, 127.8, 119.8 (1 arom. C, imidazole C(4), C(5), triazole C(3)); 129.4, 128.6, 127.6 (5 arom. CH); 126.3 (imidazole C(2)); 49.1 (PhCH<sub>2</sub>); 41.0 (CH<sub>2</sub>N); 30.7, 19.4 (2 CH<sub>2</sub>); 13.6, 9.1 (2 Me). HR-ESI-MS: 350.1589 ( $[M+\text{Na}]^+$ , C<sub>17</sub>H<sub>21</sub>N<sub>5</sub>NaO<sub>2</sub>; calc. 350.1588), 328.1771 ( $[M+\text{H}]^+$ , C<sub>17</sub>H<sub>22</sub>N<sub>5</sub>O<sub>2</sub>; calc. 328.1768).

*4-(4-Butyl-4,5-dihydro-5-oxo-1H-1,2,4-triazol-3-yl)-1,5-dimethyl-1H-imidazole 3-Oxide (8b)*. Yield: 0.176 g (70%). Colorless crystals. M.p. 214 – 216° (MeOH). IR (KBr): 3409 $vs$  (NH), 1698 $vs$  (C=O), 1559 $vs$ , 1410 $s$ , 649 $m$ .  $^1\text{H-NMR}$  ((D<sub>6</sub>)DMSO): 8.33 ( $s$ , H–C(2)); 3.71 ( $q$ ,  $J$  = 7.0, CH<sub>2</sub>N); 3.58 ( $s$ , MeN); 2.16 ( $s$ , Me); 1.64 (br.  $s$ , NH); 1.39 – 1.35, 1.10 – 1.06 ( $2m$ , 2 CH<sub>2</sub>); 0.74 ( $t$ ,  $J$  = 7.4, Me).  $^{13}\text{C-NMR}$  ((D<sub>6</sub>)DMSO): 155.6 (C=O); 137.4, 128.2, 118.8 (imidazole C(4), C(5), triazole C(3)); 126.4 (imidazole C(2)); 41.1 (CH<sub>2</sub>N); 32.7 (MeN); 32.7, 19.5 (2 CH<sub>2</sub>); 13.7, 8.9 (2 Me). HR-ESI-MS: 290.1012 ( $[M+\text{K}]^+$ , C<sub>11</sub>H<sub>17</sub>KN<sub>5</sub>O<sub>2</sub>; calc. 290.1014), 274.1273 ( $[M+\text{Na}]^+$ , C<sub>11</sub>H<sub>17</sub>N<sub>5</sub>NaO<sub>2</sub>; calc. 274.1275), 252.1451 ( $[M+\text{H}]^+$ , C<sub>11</sub>H<sub>18</sub>N<sub>5</sub>O<sub>2</sub>; calc. 252.1455).

7. *Synthesis of N-Acetylhydrazides 9. General procedure.* A mixture of hydrazide **1** (1 mmol) and Ac<sub>2</sub>O (1.1 mmol) in EtOH (5 ml) was heated to reflux for 2 h. The formed product was then filtered off, washed with Et<sub>2</sub>O, and crystallized from MeOH.

*N'-Acetyl-1-benzyl-5-methyl-3-oxido-1H-imidazolium-4-carbohydrazide (9a)*. Yield: 1.251 g (87%). Colorless crystals. M.p. 164 – 166° (MeOH). IR (KBr): 3119 $s$ , 1663 $vs$  (C=O), 1601 $vs$ , 1497 $m$ , 709 $m$ .  $^1\text{H-NMR}$  ((D<sub>6</sub>)DMSO): 12.65, 10.26 (2 br.  $s$ , 2 NH); 8.68 ( $s$ , H–C(2)); 7.41 – 7.23 ( $m$ , 5 arom. H); 5.24 ( $s$ , CH<sub>2</sub>); 2.42, 1.90 ( $2s$ , 2 Me).

$^{13}\text{C}$ -NMR ((D<sub>6</sub>)DMSO): 167.3, 159.7 (2 C=O); 135.6, 131.2, 120.8 (1 arom. C, C(4), C(5)); 129.5, 128.6, 127.6 (5 arom. CH); 126.9 (C(2)); 48.7 (CH<sub>2</sub>); 20.9, 9.6 (2 Me). HR-ESI-MS: 311.1118 ([M+Na]<sup>+</sup>, C<sub>14</sub>H<sub>16</sub>N<sub>4</sub>NaO<sub>3</sub>; calc. 311.1115); 289.1297 ([M+H]<sup>+</sup>, C<sub>14</sub>H<sub>17</sub>N<sub>4</sub>O<sub>3</sub>; calc. 289.1295).

*N'*-Acetyl-1-cyclohexyl-5-methyl-3-oxido-1H-imidazolium-4-carbohydrazide

(**9b**): Yield: 0.246 g (88%). Colorless crystals. M.p. 184–188° (MeOH). IR (KBr): 3247s (NH), 2937s, 1654vs (C=O), 1601vs, 1418m, 1047m.  $^1\text{H}$ -NMR ((D<sub>6</sub>)DMSO): 12.61 (br. s, NH); 8.68 (s, H-C(2)); 4.10–4.03 (m, CH); 2.55 (s, Me); 1.93–1.78 (m, 4 cyclohexyl H, Me); 1.66–1.59 (m, 3 cyclohexyl H); 1.43–1.36 (m, 2 cyclohexyl H); 1.20–1.13 (m, 1 cyclohexyl H).  $^{13}\text{C}$ -NMR ((D<sub>6</sub>)DMSO): 168.6, 158.0 (2 C=O); 130.7, 119.8 (C(4), C(5)); 124.5 (C(2)); 55.1 (CH); 33.0, 25.3, 24.9 (5 cyclohexyl CH<sub>2</sub>); 20.9, 9.4 (2 Me). HR-ESI-MS: 303.14267 ([M+Na]<sup>+</sup>, C<sub>13</sub>H<sub>20</sub>N<sub>4</sub>NaO<sub>3</sub>; calc. 303.1428); 281.1607 ([M+H]<sup>+</sup>, C<sub>13</sub>H<sub>21</sub>N<sub>4</sub>O<sub>3</sub>; calc. 281.1608).

8. *Synthesis of 1H-Imidazole-2-thiones 11. General Procedure.* To a magnetically stirred soln. of 1H-imidazole N-oxide **5g** or **6e** (1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 ml), a soln. of 2,2,4,4-tetramethylcyclobutane-1,3-dithione (**10**, 0.6 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 ml) was added dropwise. The addition was complete after *ca.* 10 min, and stirring was continued for 48 h, while a precipitate was formed. Then, the solvent was removed under reduced pressure, the resulting solid was washed with hexane, and filtered. The crude product was recrystallized from MeOH.

*1-Cyclohexyl-4-(4,5-dihydro-4-methyl-5-thioxo-1H-1,2,4-triazol-3-yl)-5-methyl-1H-imidazole-2(3H)-thione (11a).* Yield: 0.250 g (81%). Colorless crystals. M.p. 276 – 278° (MeOH). IR (KBr): 3092s, 2935s, 1560m, 1500m, 1416m, 1348m.  $^1\text{H}$ -NMR ((D<sub>6</sub>)DMSO): 3.72 (s, MeN); 3.35 – 3.29 (m, CH); 2.33 (s, Me); 1.83 – 1.64 (m, 6 cyclohexyl H); 1.36 – 1.16 (m, 4 cyclohexyl H); NH absorption missing.  $^{13}\text{C}$ -NMR

((D<sub>6</sub>)DMSO): 171.6, 163.2 (2 C=S); 141.2 (triazole C(3)); 126.3, 116.5 (imidazole C(4), C(5)); 55.4 (CH); 31.7 (MeN); 26.3, 25.8, 21.9 (5 cyclohexyl CH<sub>2</sub>); 11.9 (Me). HR-ESI-MS: 332.0973 ([M+Na]<sup>+</sup>, C<sub>13</sub>H<sub>19</sub>N<sub>5</sub>NaS<sub>2</sub>; calc. 332.0974), 310.1150 ([M+H]<sup>+</sup>, C<sub>13</sub>H<sub>20</sub>N<sub>5</sub>S<sub>2</sub>; calc. 310.1155).

*1-Cyclohexyl-5-methyl-4-[5-(4-sulfophenyl)amino]-1,3,4-thiadiazol-2-yl)-1H-imidazole-2(3H)-thione (11b)*. Yield: 0.338 g (91%). Colorless crystals. M.p. 314 – 316° (dec., MeOH). IR (KBr): 3400–2700*m* (br), 3178*s*, 2935*s*, 1405*s* (br), 1122*s* (br), 616*m*. <sup>1</sup>H-NMR ((D<sub>6</sub>)DMSO): 7.57 (br. *s*, 4 arom. H); 3.36 – 3.29 (*m*, CH); 2.57 (*s*, Me); 1.94 – 1.63 (*m*, 6 cyclohexyl H); 1.37 – 1.21 (*m*, 4 cyclohexyl H); NH absorption missing. <sup>13</sup>C-NMR ((D<sub>6</sub>)DMSO): 163.9, 147.3 (thiadiazole C(2), C(5)); 162.9 (C=S); 142.9, 123.0, 120.2, 116.8 (2 arom. C, imidazole C(4), C(5)); 140.1, 127.1 (4 arom. CH); 57.1 (CH); 26.3, 25.3, 21.2 (5 cyclohexyl CH<sub>2</sub>); 11.9 (Me). HR-ESI-MS: 474.0695 ([M+Na]<sup>+</sup>, C<sub>18</sub>H<sub>21</sub>N<sub>5</sub>NaO<sub>3</sub>S<sub>3</sub>; calc. 474.0699), 452.0877 ([M+H]<sup>+</sup>, C<sub>18</sub>H<sub>22</sub>N<sub>5</sub>O<sub>3</sub>S<sub>3</sub>; calc. 452.0879).

## REFERENCES

- [1] C. A. M. Fraga, E. J. Barreiro, *Curr. Med. Chem.* **2006**, *13*, 167; S. Rollas, Ş. G. Küçükgülzel, *Molecules* **2007**, *12*, 1910; B. Narasimhan, P. Kumar, D. Sharma, *Acta Pharm. Sci.* **2010**, *52*, 169.
- [2] T. P. Sycheva, T. N. Pavlova, M.N. Shchukina, *Pharm. Chem. J.* **1972**, *6*, 696.
- [3] J. H. M. Lange, H. H. Van Stuivenberg, H. K. A. C. Hein, T. J. P. Adolfs, A. C. McCreary, H. G. Keizer, H. C. Wals, W. Veerman, A. J. M., W. De Looft, P. C. Verveer, and C. G. Kruse, *J. Med.Chem.* **2005**, *48*, 1823.


- [4] Takeda Chemical Industries, Ltd., 1998, US 1998/5753664 A1; Solvay Pharmaceuticals B. V., 2005, US 2005/54679 A1; Astrazeneca UK Ltd., 2007, WO 2007/31721 A1.
- [5] A. A. Hassan, A. M. Shawky, *J. Heterocycl. Chem.* **2010**, *47*, 745.
- [6] G. Mlostoń, A. Pieczonka, E. Kowalczyk, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **2011**, *94*, 1764.
- [7] M. Jasiński, G. Mlostoń, A. Linden, H. Heimgartner, *Helv. Chim. Acta* **2008**, *91*, 1916; M. Jasiński, G. Mlostoń. *Coll. Czech. Chem. Commun.* **2010**, *75*, 871.
- [8] a) G. Mlostoń, T. Gendek, H. Heimgartner, *Helv. Chim. Acta* **1998**, *81*, 1585; b) G. Mlostoń, M. Celeda, G. K. S. Prakash, G. A. Olah, H. Heimgartner, *Helv. Chim. Acta* **2000**, *83*, 728; c) G. Mlostoń, J. Romański, M. Jasiński, H. Heimgartner, *Tetrahedron: Asymmetry* **2009**, *20*, 1073; d) G. Mlostoń, M. Jasiński, H. Heimgartner, *Eur. J. Org. Chem.* **2011**, 2542; e) G. Mlostoń, M. Jasinski, D. Rygielska, H. Heimgartner, *Heterocycles* **2011**, *83*, 765; f) G. Mlostoń, D. Rygielska, M. Jasiński, H. Heimgartner, *Tetrahedron: Asymmetry* **2011**, *22*, 669.
- [9] S. Hauptmann, *Organische Chemie*, 2. Aufl., VEB Deutscher Verlag für Grundstoffindustrie, Leipzig, 1985, p. 511.
- [10] R. N. Khelevin, *Zh. Org. Khim.* **1984**, *20*, 379 and 1290; R. N. Khelevin, *Zh. Obsh. Khim.* **1986**, *56*, 202 and 2315.
- [11] I. P. Singh Kapoor, M. Kapoor, G. Singh, *J. Therm. Anal. Calorim.* **2010**, *102*, 723.
- [12] X. J. Zou, G. Y. Jin, Z. X. Zhang, *J. Agricul. Food Chem.* **2002**, *50*, 1451; A. Maliszewska-Guz, M. Wujec, M. Pitucha, M. Dobosz, A. Chodkowska, E.

Jagiello-Wójtowicz, L. Mazur, A. E. Kozioł, *Collect. Czech. Chem. Commun.* **2005**, *70*, 51.


- [13] A. P. Liesen, T. M. de Aguiño, C. S. Carvalho, V. T. Lima, J. M. de Araújo, J. G. de Lima, A. R. de Faria, E. J. T. de Melo, A. J. Alves, E. W. Alves, A. Q. Alves, A. J. S. Góes, *Eur. J. Med. Chem.* **2010**, *45*, 3685.




Scheme 1




Scheme 2




Scheme 3



Scheme 4



## Scheme 5



## Graphical Abstract

